

PA-058

BP2202: An Off-the-Shelf iPSC-Derived BCMA-Targeted Cellular Therapy for Multiple Myeloma



K. Shioya¹, A. Urakami¹, T. Shigeura¹, N. Watanabe¹, M. Fujimaki¹, YC. Lin¹, N. Komatsuzaki¹, T. Matsumura¹, K. Nakajima¹, R. Kuniyoshi¹, Y. Ohira¹, S. Motohashi², H. Koseki³, L. Zhang¹

¹ BrightPath Biotherapeutics Co., Ltd.; Tokyo, Japan, ² Department of Medical Immunology, Graduate School of Medicine, Chiba University; Chiba, Japan, ³ Laboratory for Developmental Genetics, RIKEN Center for Integrative Medical Sciences; Yokohama, Japan.

INTRODUCTION

BCMA-targeted CAR-T cells and T-cell engagers have shown meaningful activity in relapsed/refractory multiple myeloma (RRMM), but relapse remains common. Autologous CAR-T therapy is limited by manufacturing complexity, long vein-to-vein time, and product variability, while conventional allogeneic CAR-T approaches have shown limited persistence. Thus, off-the-shelf allogeneic therapies with durable efficacy are needed.

Invariant natural killer T (iNKT) cells are a rare T-cell subset with potent direct and indirect anti-tumor activity through activation of CD8+ T cells and other immune populations. iPSC technology enables scalable generation of this otherwise rare cell population while preserving its functional properties.

BP2202 is an off-the-shelf, iPSC-derived BCMA CAR-iNKT cell therapy that leverages both CAR-mediated cytotoxicity and the intrinsic anti-tumor functions of iNKT cells, while enabling scalable, master cell bank-based manufacturing.

AIM

- 1. *In vitro* characterization**
Anti-tumor activity and functional properties against BCMA-expressing myeloma cells
- 2. *In vivo* efficacy**
Anti-tumor activity
- 3. Clinical translation**
Design of a Phase 1 study evaluating the safety and preliminary efficacy of BP2202 in RRMM

METHOD

- BP2202 was evaluated for phenotype, CAR expression, antigen-specific cytotoxicity, and cytokine production.
- Cytotoxicity was assessed by luciferase-based killing assay (RPMI8226, BCMA-positive; SK-OV-3, BCMA-negative) and by xCELLigence assay (U-2OS, CD1d-positive, $\pm$ α -GalCer). Anti-tumor activity was evaluated in MM.1S xenograft mouse models by bioluminescence imaging.
- Genomic safety of the gene-edited iPSC was assessed by karyotyping, optical genome mapping, whole genome sequencing, and off-target analysis.
- The phase 1 study is an open-label, multicenter, dose-escalation trial in RRMM patients after ≥ 3 prior lines of therapy.

RESULTS

1. Overview of BP2202 manufacturing




Figure 1. iNKT cells were reprogrammed into iPSCs, followed by insertion of the BCMA-CAR and IL-15 genes into the iPSC genome using gene-editing technology. After single-cell cloning and screening, a CAR-iPSC clone was established and used to generate the master cell bank (MCB). The MCB underwent extensive quality control, including genomic safety assessment. Manufacturing of BP2202 begins with iPSC expansion, followed by primary differentiation into Hematopoietic Stem and Progenitor Cells (HSPCs) and subsequent differentiation into BP2202.

3. Phenotype and CAR expression




Figure 2. Surface marker expression of BP2202 by Flow cytometry. Histograms show the expression of CAR and HLA molecules (HLA-ABC and HLA-DP/DR/DQ) in the live cell population. The expression of the iNKT TCR (Va24-Ja18), CD3, CD4, and CD8a, as well as the NK-associated markers CD56, NKG2A, NKG2D, DNAM-1, Nkp30, Nkp44, and Nkp46, was analyzed.

4. Killing activity and cytokine production






Figure 3. (A) BP2202, with or without 100 ng/mL α -GalCer, was co-cultured with 1 $\times$ 10⁴ U-2OS cells/well at various E/T ratios for 72h, and cytotoxic activity was evaluated using the real-time cell analyzer xCELLigence. (B) BP2202 was co-cultured with 3 $\times$ 10⁴ target cells/well at various E/T ratios for 24h, and cytotoxic activity was evaluated using a luciferase assay. Cytotoxicity was calculated as follows: % cytotoxicity = [1 - (RFU of co-cultured cells/RFU of target cells alone)] $\times$ 100. Data are presented as mean $\pm$ SEM (n = 3). (C) Expression pattern of BCMA, CD1d and NK receptors ligands using flowcytometry analysis. NA: Not Assessed (D) BP2202 was co-cultured with 1 $\times$ 10⁵ target cells/well at an E/T ratio of 1 for 24 h, and IFN- γ and TNF- α levels in the culture supernatant were measured using a Cytometric Bead Array. Data are presented as mean $\pm$ SEM (n = 3).

2. Key features of BP2202





- BCMA-CAR Knock-in**
 $\Rightarrow$ Enables targeted killing of BCMA⁺ tumor cells
- Soluble hIL-15 expression**
 $\Rightarrow$ Enhances iNKT cell survival and persistence
- HLA-class I Knock-out**
 $\Rightarrow$ Reduces immunogenicity and promotes persistence
- Invariant TCR expression**
 $\Rightarrow$ Enables CD1d-restricted cytotoxicity, Promotes DC maturation, Facilitates M1 ϕ polarization
- NK receptor expression**
 $\Rightarrow$ Boost innate cytotoxicity

5. *In vivo* anti-tumor effect



6. Study design for Phase 1 clinical trial (BP2202-001)



CONCLUSIONS

- BP2202 retains key iNKT characteristics, including NK receptors and invariant TCR expression, supporting multi-MOA activity.
- BP2202 mediates anti-tumor activity through CAR-, NK receptor-, and CD1d/TCR pathways.
- No unintended structural or genomic abnormalities or off-target events were detected.
- BP2202 demonstrated potent anti-tumor activity in vitro and significant tumor growth inhibition in vivo.
- These findings support the clinical development of BP2202 as a promising off-the-shelf therapy for R/R MM.

CONTACT INFORMATION

Lilin Zhang, MD, PhD.,
Director, Pharmaceutical Development
BrightPath Biotherapeutics Co., Ltd
zhang_l@brightpathbio.com

